

Quantum clocks, mirrors and Alice and Bob in Gravity

Časlav Brukner



Olomouc, May 10th, 2012

Motivation

Quantum Mechanics

- entanglement
- single particle interference
- Bohr's complementarity principle
- Born's rule



Passed

Newtonian gravity sufficient
(if any gravity effects seen at all!)

General Relativity

- Einstein's equations
- gravity as space-time geometry
- gravitational time dilation
- black holes



Passed

consistent with classical
mechanics

Motivation

Quantum Mechanics

General Relativity


1. Effects that require both theories to be explained?
2. Effects that require an unified framework („quantum gravity“)?



Outline


- Introduction & motivation
- [1] Gravitational redshift and quantum complementarity
- [2] Quantum correlations with no-causal order
- [3] Probing Planck-scale physics with quantum optics
- Conclusion

[1] Gravitational redshift and quantum complementarity




M. Zych, F. Costa, I. Pikovski, Č. Brukner:
Nature Communication 2:505
doi: 10.1038/ncomms1498 (2011)

Quantum Complementarity Principle




Quantum Complementarity Principle




It is **not possible** to simultaneously know the path of the particle and observe its interference.

Gravitational time dilation



Two initially synchronized clocks placed at different gravitational potentials.

Clock closer to a massive body ticks slower than the clock further away from the mass.




Initially synchronized clocks will eventually show **different times** when placed at different gravitational potentials.

Interference of clocks


GR: time shown by the clock depends on the path taken

running clock in a superposition



QM: interference cannot be observed if path can be known


switched off clock
in a superposition



interference can
be observed


quantum complementarity + time dilation =
= drop in the interferometric visibility

Mach-Zehnder interferometer in a gravitational field



$\gamma_{1,2}$: two possible paths through the setup,
 g : homogeneous gravitational field,
 Δh : separation between the paths

Quantum Complementarity



“clock” - a system with an evolving in time degree of freedom

modes associated with the path γ_1

state of the “clock”, which followed path γ_1

$$|\Psi_{MZ}\rangle = \frac{1}{\sqrt{2}} (i|r_1\rangle|\tau_1\rangle e^{-i\phi_1} + |r_2\rangle|\tau_2\rangle e^{-i\phi_2 + i\varphi})$$

Probabilities of detection

$$P_{\pm} = \frac{1}{2} \pm \frac{1}{2} |\langle\tau_1|\tau_2\rangle| \cos(\Delta\phi + \alpha + \varphi)$$

$$\langle\tau_1|\tau_2\rangle = |\langle\tau_1|\tau_2\rangle| e^{i\alpha}$$

Visibility of the interference pattern:

$$\mathcal{V} = |\langle\tau_1|\tau_2\rangle|$$

Distunguishability of the paths:

$$\mathcal{D} = \sqrt{1 - |\langle\tau_1|\tau_2\rangle|^2}$$

Interferometric visibility drops to the extent to which path information becomes available from the “clock”

Results

$$H_{\odot} = E_0 |0\rangle\langle 0| + E_1 |1\rangle\langle 1|$$

$$|\tau^{in}\rangle = \frac{1}{\sqrt{2}} (|0\rangle - |1\rangle)$$

- $\Delta E := E_1 - E_0$
- $\Delta V := g\Delta h$, gravitational potential
- Δh : distance between the paths
- ΔT : time for which the particle travels in superposition at constant heights

$$P_+(\varphi, m, \Delta E, \Delta V, \Delta T) =$$

$$= \frac{1}{2} \pm \frac{1}{2} \cos\left(\frac{\Delta E \Delta V \Delta T}{2\hbar c^2}\right) \cos\left(\left(mc^2 - \langle H_{\odot} \rangle + \bar{E}_{GR}^{corr}\right) \frac{\Delta V \Delta T}{\hbar c^2} + \varphi\right)$$

relative phase from the Newtonian potential

GR corrections to the relative phase from the path d.o.f.

new effects appearing with the "clock":

change in the interferometric visibility

$$\mathcal{V} = \left| \cos\left(\frac{\Delta E \Delta V \Delta T}{2\hbar c^2}\right) \right|$$

phase shift proportional to the average internal energy

Results


$$H_{\odot} = E_0 |0\rangle\langle 0| + E_1 |1\rangle\langle 1|$$

$$|\tau^{in}\rangle = \frac{1}{\sqrt{2}} (|0\rangle - |1\rangle)$$

- $\Delta E := E_1 - E_0$
- $\Delta V := g\Delta h$, gravitational potential
- Δh : distance between the paths
- ΔT : time for which the particle travels in superposition at constant heights

$$P_+(\varphi, m, \Delta E, \Delta V, \Delta T) =$$

$$= \frac{1}{2} \pm \frac{1}{2} \cos\left(\frac{\Delta E \Delta V \Delta T}{2\hbar c^2}\right) \cos\left((mc^2 - \langle H_{\odot} \rangle + \bar{E}_{corr}) \frac{\Delta V \Delta T}{\hbar c^2} + \varphi\right)$$



- dashed, black line - interference with the “clock” switched off
- blue line - phase with the “clock” switched on
- thick, red line - modulation in the visibility

Phase shift vs Drop of visibility

$$|\Psi_{MZ}\rangle = \frac{1}{\sqrt{2}} (i|r_1\rangle|\tau_1\rangle e^{-i\phi_1} + |r_2\rangle|\tau_2\rangle e^{-i\phi_2 + i\varphi})$$

Phase Shift

Explainable by:

- a potential force in absolute time (possible non-Newtonian)
- analogue to a charged particle in EM field
- Flat space-time: no redshift
- independent of whether a particle is a „clock“ or a rock

Colella, R., Overhauser, A. W. & Werner, S. A. *Phys. Rev. Lett.* 34, 1472–1474 (1975).

Müller, H., Peters, A. & Chu, S. *Nature* 463, 926–929 (2010).

Drop in Visibility

Not explainable without:

- gravity as metric theory,
- proper time τ flows at different rates – redshift
- curved space-time geometry
- iff a particle is an operationally well defined „clock“

Experiment challenging (2-3 orders of magnitude)


Snímek 15

MZ3

such an interpretation was recently proposed in: H. Müller, A. Peters, & S. Chu, A precision measurement of the gravitational redshift by the interference of matter waves. *Nature* 463, 926–929 (2010).


Magdalena Zych; 23.1.2012

[2] Quantum correlations with no causal order




O. Oreshkov, F. Costa, Č. Brukner:
arXiv:1105.4464

Measurements in space-time



- Fix positions
- Define initial state
- Follow Eqs of motion
- Include causal influences
- Find joint probabilities $P(A, B, C, D, E)$


Circuit model



Space-time & definite causal structure are pre-existing entities.


What happens if one removes global time and causal structure from quantum mechanics? What new phenomenology is implied?

Measurements in space-time




- Fix positions
- Define initial state
- Follow Eqs of motion
- Include causal influences
- Find joint probabilities $P(A, B, C, D, E)$

Circuit model



New computational model?
New phenomenology?

Operational approach



The system exits the lab.




One out of a set of possible transformations (CP-maps) is performed.

A system enters the lab.

This is the **only** way how the labs interact with the “outside world”.

Operational approach


„Process“
(„catalogue of our knowledge“)



No prior assumption of pre-existing causal structure, in particular of the pre-existing background time.

Main premise:


Local quantum mechanics: The local operations of each party are described by quantum mechanics.




Transformations = **completely positive** (CP)
trace non increasing maps

$$\mathcal{M}_j: \mathcal{L}(\mathcal{H}^2) \rightarrow \mathcal{L}(\mathcal{H}^1)$$

Two parties



$$\mathcal{M}_{j_A}^A : \mathcal{L}(\mathcal{H}^{A_2}) \rightarrow \mathcal{L}(\mathcal{H}^{A_1})$$




$$\mathcal{M}_{j_B}^B : \mathcal{L}(\mathcal{H}^{B_2}) \rightarrow \mathcal{L}(\mathcal{H}^{B_1})$$

Probabilities are **bilinear** functions of the CP maps

Choi-Jamilkowski representation of the CP maps:


$$P(\mathcal{M}^A, \mathcal{M}^B) = \text{Tr} \left[\underbrace{W^{A_1 A_2 B_1 B_2}}_{\text{„Process Matrix“}} \left(\underbrace{\rho_{\mathcal{M}^A}^{A_1 A_2}}_{\text{CP maps}} \otimes \underbrace{\rho_{\mathcal{M}^B}^{B_1 B_2}}_{\text{CP maps}} \right) \right]$$

Bipartite state




Sharing a joint state; No signalling

Channel $B \rightarrow A$



Sending a state from B to A; Possibility of signalling

Channel $A \rightarrow B$



Sending a state from A to B; Possibility of signalling

Mixtures of different orders also possible

Most general causally separable situation:
probabilistic mixture of ordered ones:

$$W^{A_1 A_2 B_1 B_2} = q W^{B \not\prec A} + (1 - q) W^{A \not\prec B}$$

↑
Signalling only from A
to B or causally
independent


↑
Signalling only from B
to A or causally
independent

Do all possible processes W respect definite
causal order?

NO!

Causal Game

Guess **x** is produced
in **local time before**
bit **a** is given




- Alice is given bit **a** and Bob bit **b**.
- Alice produces **x** and Bob **y**, which are their best guesses for the value of the bit given to the other.
- Bob is given an additional bit **b'** that tells him whether he should guess her bit (**b'=1**) or she should guess his bit (**b'=0**).
- The goal is to maximize the probability for correct guess:

$$p_{succ} := \frac{1}{2} [P(x = b|b' = 0) + P(y = a|b' = 1)]$$

Causally ordered situation

Case: $B \not\prec A$

Global Time




$$P(y = a | b' = 1) = 1$$

$$P(x = b | b' = 0) = 1/2$$

$$p_{succ} = P(x = b | b' = 0) + P(y = a | b' = 1) \leq \frac{3}{4}$$

Causally non-separable situation

$$W^{A_1 A_2 B_1 B_2} = \frac{1}{4} \left[\mathbb{1} + \frac{1}{\sqrt{2}} (\sigma_z^{A_1} \sigma_z^{B_2} + \sigma_z^{A_2} \sigma_z^{B_1} \sigma_x^{B_2}) \right]$$



"I think you should be more explicit here in step two."

The probability of success is

$$p_{succ} = \frac{2+\sqrt{2}}{4} > \frac{3}{4}$$

"Tsirlason bound for non-causal correlations" ??

This process cannot be realized as a probabilistic mixture of causally ordered situations!

[2] Probing Planck physics with quantum optics



I. Pikovski, M. R. Vanner, M. Aspelmeyer, M. S. Kim and Č. Brukner:
Nature Physics (2012) doi:10.1038/nphys2262

Experimental quantum gravity?

Effects largely believed to be relevant at the **Planck-scale**:

$$E_{Planck} = \sqrt{\frac{\hbar c^5}{G}} = 1.956 \times 10^9 \text{ J}$$

$$\frac{E_{exp}}{E_{Planck}} \approx 10^{-15}$$

High-energy scattering experiments

$$L_{Planck} = \sqrt{\frac{\hbar G}{c^3}} = 1.6161 \times 10^{-35} \text{ m}$$

$$\frac{x_{exp}}{L_{Planck}} \approx 10^{17}$$

High-precision quantum metrology

$$m_{Planck} = \sqrt{\frac{\hbar c}{G}} = 22 \text{ } \mu\text{g}$$

Optomechanics:

Mirrors can have mass of pg – kg!

Modified uncertainty relation

- A minimal measurable length scale $\Delta x_{min} \sim$ Planck-length $L_p \sim 10^{-35} m$.
- Thus $\Delta x \Delta p = \frac{\hbar}{2}$ cannot hold for $\Delta p \rightarrow \infty$
- Typical modification in QGR:

$$\Delta x \Delta p \geq \frac{\hbar}{2} \left(1 + \beta_0 \frac{\Delta p^2}{M_{Pl}^2 c^2} \right)$$


standard QM

(L. Garay, *Int. J. Mod. Phys. A10*, 145 (1995))

Modification: $M_{Pl} \approx 22 \mu g$
Planck-mass, β_0 dimensionless parameter

Current experimental bound: $\beta_0 < 10^{33}$

(S. Das & E. C. Vagenas, *PRL* 101, 221301 (2008))



Possible commutator modifications

$\Delta x \Delta p \geq \frac{\hbar}{2} (1 + \beta \Delta p^2)$ implies a modified commutator. E.g.:

- $[\hat{X}, \hat{P}]_{\beta} = i \left(1 + \beta_0 \frac{\hat{P}^2}{M_{Pl}^2 c^2} \right)$ *(A. Kempf, G. Mangano and R. Mann, PRD, 52, 2 (1995))*
- $[\hat{X}, \hat{P}]_{\mu} = i \sqrt{1 + 2\mu_0 \frac{(\hat{P}/c)^2 + m^2}{M_{Pl}^2}}$ *(M. Maggiore, Phys. Lett. B, 319 (1993))*
- $[\hat{X}, \hat{P}]_{\gamma} = i \left(1 - \gamma_0 \frac{\hat{P}}{M_{Pl} c} + \gamma_0^2 \frac{\hat{P}^2}{M_{Pl}^2 c^2} \right)$ *(A. F. Ali, S. Das and E. C. Vagenas, Phys. Lett. B, 678 (2009))*


Note: ground-state $p_0 = \sqrt{m\omega\hbar}$, mass-dependent

Opto-mechanics

Control of a massive systems with light

Opto-mechanical interaction:

$$\hat{H} = \hbar\omega_m \hat{n}_m + \hbar\omega_0 \hat{n}_L - \hbar g_0 \hat{n}_L \hat{X}_m$$




$$g_0 = \frac{\omega_0}{L} \sqrt{\frac{\hbar}{m\omega_m}} \text{ Opto-mechanical coupling rate}$$

- Pulsed interactions (duration $\tau \ll \omega_m$): *(Vanner, et al., PNAS 108, 16182 (2011))*

$$\hat{H} \approx -\hbar g_0 \hat{n}_L \hat{X}_m$$

- Harmonic evolution: $\hat{X}_m(t) = \hat{X}_m \cos(\omega_m t) - \hat{P}_m \sin(\omega_m t)$




Loop in a phase space

Displacements of a quantum system around a loop in phase space via an ancillary (light) system:

$$\begin{aligned}\hat{\xi} &= e^{i\lambda\hat{n}_L\hat{P}} e^{-i\lambda\hat{n}_L\hat{X}} e^{-i\lambda\hat{n}_L\hat{P}} e^{i\lambda\hat{n}_L\hat{X}} \\ &= e^{-i\lambda^2\hat{n}_L^2}\end{aligned}$$

Four pulses separated by $\omega_m t = \pi/2$:

- Resulting phase changes the ancilla, but is state-independent
- Mechanics remains unaffected, and is fully disentangled from the ancilla



Phase due to QG


$$\hat{\xi} = e^{i\lambda\hat{n}_L\hat{P}} e^{-i\lambda\hat{n}_L\hat{X}} e^{-i\lambda\hat{n}_L\hat{P}} e^{i\lambda\hat{n}_L\hat{X}} = e^{\sum_{k=1}^{\infty} \frac{(-i\lambda\hat{n}_L)^{k+1}}{k!} [\hat{X}, \hat{P}]_k}$$

where $[\hat{X}, \hat{P}]_k \equiv [\hat{X}, [\hat{X}, \dots, \hat{P}]]_k$

- QM: $[\hat{X}, \hat{P}] = i \Rightarrow \hat{\xi}_{QM} = e^{-i\lambda^2\hat{n}_L^2}$
- QG: $[\hat{X}, \hat{P}] = i F(\hat{X}, \hat{P})$

Any arbitrary deformed algebra will show in $\hat{\xi}$!

By measuring the ancilla (initially in $|\alpha\rangle$) one can obtain a measure of the commutator.



$$\langle \hat{a}_L \rangle = \langle \alpha | \hat{\xi}^\dagger \hat{a}_L \hat{\xi} | \alpha \rangle \cong \langle \hat{a}_L \rangle_{QM} e^{-i \Theta([\hat{X}, \hat{P}]_{mod})}$$

Example: $\Theta(\beta) \simeq \frac{4}{3} \beta N_p^3 \lambda^4 e^{-i6\lambda^2}$

Table 2 | Experimental parameters to measure quantum gravitational deformations of the canonical commutator.


	$[X_m, P_m]$	Equation (2)	Equation (3)	Equation (1)
	$ \Theta $	$\mu_0 \frac{32\hbar \mathcal{F}^2 m N_p}{M_p^2 \lambda_L^2 \omega_m}$	$\gamma_0 \frac{96\hbar^2 \mathcal{F}^3 N_p^2}{M_p c \lambda_L^3 m \omega_m}$	$\beta_0 \frac{1024\hbar^3 \mathcal{F}^4 N_p^3}{3M_p^2 c^2 \lambda_L^4 m \omega_m}$
Finesse	\mathcal{F}	10^5	2×10^5	4×10^5
Mass	m	10^{-11} kg	10^{-9} kg	10^{-7} kg
Mech. Frequency	$\omega_m/2\pi$	10^5 Hz	10^5 Hz	10^5 Hz
Optical wavelength	λ_L	1,064 nm	1,064 nm	532 nm
Photon number	N_p	10^8	5×10^{10}	10^{14}
Measurement runs	N_r	1	10^5	10^6
Measur. precision	$\delta\langle\Phi\rangle$	10^{-4}	10^{-8}	10^{-10}

The parameters are chosen such that a precision of $\delta\mu_0 \sim 1$, $\delta\gamma_0 \sim 1$ and $\delta\beta_0 \sim 1$ can be achieved, which amounts to measuring Planck-scale deformations.

Quantum Information Meets Gravity


Summary

1. New paradigm for tests of genuine general relativistic effects in quantum mechanics:
 - **Drop in the visibility of quantum interference due to gravitational time dilation**
2. Quantum formalism for indefinite causal structures
 - **Quantum correlations with no-causal order**
3. Possibility to probe phenomenological predictions of quantum gravity in massive quantum systems:
 - **Measurement of the deformation of commutation relation of the center-of-mass modes**



Borivoje Dakic

Igor Pikovski



Fabio Costa

Magdalena Zych



Ognyan Oreshkov (U. Brussels)

C.B.



CoQUS Complex Quantum Systems

FOXi John Templeton Foundation
 FOUNDATIONAL QUESTIONS INSTITUTE

Thank you for your attention

quantumfoundations.weebly.com